

# The Relevance of *In Silico*, *In Vitro* and Non-human Primate Based Approaches to Clinical Research on Major Depressive Disorder

Alternatives to Laboratory Animals  
2019, Vol. 47(3-4) 128–139  
© The Author(s) 2019



Article reuse guidelines:  
sagepub.com/journals-permissions  
DOI: 10.1177/0261192919885578  
journals.sagepub.com/home/atl



Constança Carvalho<sup>1</sup>, Susana A.M. Varela<sup>2,3</sup>,  
Luísa Ferreira Bastos<sup>4,5</sup>, Inês Orfão<sup>1,2</sup>, Vanda Beja<sup>6</sup>,  
Manuel Sapage<sup>2</sup>, Tiago A. Marques<sup>7,8</sup>, Andrew Knight<sup>9</sup>  
and Luís Vicente<sup>1</sup>

## Abstract

Major depressive disorder (MDD) is the most severe form of depression and the leading cause of disability worldwide. When considering research approaches aimed at understanding MDD, it is important that their effectiveness is evaluated. Here, we assessed the effectiveness of original studies on MDD by rating their contributions to subsequent medical papers on the subject, and we compared the respective contribution of findings from non-human primate (NHP) studies and from human-based *in vitro* or *in silico* research approaches. For each publication, we conducted a quantitative citation analysis and a systematic qualitative analysis of the citations. In the majority of cases, human-based research approaches (both *in silico* and *in vitro*) received more citations in subsequent human research papers than did NHP studies. In addition, the human-based approaches were considered to be more relevant to the hypotheses and/or to the methods featured in the citing papers. The results of this study suggest that studies based on *in silico* and *in vitro* approaches are taken into account by medical researchers more often than are NHP-based approaches. In addition, these human-based approaches are usually cheaper and less ethically contentious than NHP studies. Therefore, we suggest that the traditional animal-based approach for testing medical hypotheses should be revised, and more opportunities created for further developing human-relevant innovative techniques.

## Keywords

animal use alternatives, *in silico*, *in vitro*, major depressive disorder, non-human primate, three Rs

## Introduction

According to the World Health Organization, depression is the leading cause of morbidity worldwide. It affects more than 300 million people of all ages and is a major contributor to the overall global burden of disease.<sup>1</sup> People who suffer from depression are more prone to an early death either by suicide or through the development of other conditions such as cancer, heart disease or stroke.<sup>2,3</sup> In addition, these patients are also more prone to a number of other disorders (e.g. osteoporosis)<sup>4</sup> that, although not life-threatening, do significantly impact not only quality of life but also public health and national economies.

Accordingly, major investment has been dedicated to research aiming to improve the understanding of all eight forms of depression.<sup>5</sup> Major depressive disorder (MDD) is the most severe type and the third leading cause

<sup>1</sup> Centro de Filosofia das Ciências da Universidade de Lisboa, Lisboa, Portugal

<sup>2</sup> cE3c—Centre for Ecology, Evolution and Environmental Changes, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal

<sup>3</sup> Instituto Gulbenkian de Ciência (IGC), Oeiras, Portugal

<sup>4</sup> Instituto de Investigação e Inovação em Saúde (i3S), University of Porto, Porto, Portugal

<sup>5</sup> Instituto de Engenharia Biomédica (INEB), Porto, Portugal

<sup>6</sup> Independent Consultant, Clinical Psychologist, Lisboa, Portugal

<sup>7</sup> Centre for Research into Ecological and Environmental Modelling, University of St Andrews, St Andrews, Scotland, UK

<sup>8</sup> Departamento de Biologia Animal, Centro de Estatística e Aplicações, Faculdade de Ciências, Universidade de Lisboa, Lisboa, Portugal

<sup>9</sup> Centre for Animal Welfare, University of Winchester, Winchester, UK

## Corresponding author:

Constança Carvalho, Centro de Filosofia das Ciências da Universidade de Lisboa, Campo Grande 016, 1749-016 Lisboa, Portugal.

Email: constanca.carvalho@sapo.pt

of long-term disability.<sup>6</sup> Besides, the few studies that have comprehensively investigated the impact of MDD in Europe (from 2004 to 2010) have shown that MDD was the costliest brain disorder in Europe, accounting for at least 1% of the total European economy.<sup>7,8</sup> In the United States, the economic burden of MDD alone was US\$210.5 billion in 2010.<sup>9</sup>

Clinical research is expensive, time-consuming and potentially ethically contentious. For instance, every patient who enrolls in a clinical trial is subject to an increased level of risk with respect to deviations from their regular clinical care, particularly with regard to the occurrence of unexpected effects from exposure to a new treatment. Non-clinical (i.e. preclinical) research, often involving non-human animals and human-based *in vitro* and *in silico* approaches, is sometimes valuable in the early steps of biomedical research to simplify and accelerate drug and treatment discovery. However, to optimise the outcomes of this non-clinical research, it is crucial to evaluate the research approaches that might have the most potential for patient treatment results.

Animal-based research has been accepted as the ‘gold standard’ approach for preclinical biomedical research and testing since the second half of the 20th century.<sup>10</sup> Within this approach, non-human primate (NHP) research has been considered particularly relevant, due to the similarity between humans and NHPs. However, this similarity has led to NHPs being afforded various degrees of legal protection in different regions of the world. For example, Europe,<sup>11</sup> the United States<sup>12</sup> and New Zealand<sup>13</sup> have imposed considerable restrictions to the use of NHPs for scientific purposes. These restrictions are due to the understanding that subjecting NHPs to laboratory confinement alone, even before considering the use of any invasive or intrusive procedures, has resulted in psychosomatic injury, mutilation and physiological traits that have been compared to those exhibited by people with post-traumatic stress disorder.<sup>13–20</sup> Moreover, NHPs are expensive to acquire<sup>21</sup> and are the most expensive animals to maintain.<sup>22</sup>

The legislation on animal use for experimental purposes of several countries (e.g. *Directive 2010/63/EU*) requires a cost–benefit assessment to be carried out prior to conducting a procedure on a non-human animal. For each project, the likely harm to the animal should be balanced against the potential benefits, and the project should only go ahead if the expected benefits outweigh the harms inflicted to the animals involved.

Considering all of the above, it is assumed that when research is conducted on NHPs, due to the ethical and economic concerns surrounding this practice, this research should provide highly relevant data that lead to concrete improvements in patient outcomes. While some authors assert that animal research approaches, and those involving NHPs in particular, are crucial for biomedical progress,<sup>23</sup> an increasing number of evidence-based papers show that

the contribution of animal-based research to the advancement of human healthcare has been poor,<sup>24</sup> including in the case of MDD.<sup>25</sup> However, it is yet to be established whether this poor contribution is due to the intrinsic limitations of all non-clinical research, or whether human-based (*in vitro* and *in silico*) non-clinical research approaches are more effective in helping biomedical progress, at least when seeking to understand complex disorders of a multifactorial origin, such as MDD.

*In vitro* and *in silico* methods that directly rely on human-based knowledge and/or material are thought to potentially allow for faster development of medical treatments.<sup>26,27</sup> Usually, they are also more cost-effective than animal-based methods. However, despite yielding data of sufficient value to further disease understanding in humans, and providing the means to test new therapies, such non-animal methods are still judged against the standard biomedical research paradigm. Indeed, they are seen as incomplete on their own and considered to be preliminary steps prior to (often contradictory) animal testing.<sup>28,29</sup>

To shed light on this debate, the current study examines and compares the contribution of results from NHP studies, as well as from *in silico*-based and *in vitro*-based approaches, to clinical studies on MDD. This allows us to: (a) evaluate whether the low transferability of knowledge to clinical research is a common trait of all non-clinical research approaches; and (b) evaluate the specific relevance of NHP studies and human-based *in silico* and *in vitro* approaches to human clinical studies.

Considering the dominance of NHP studies within the current preclinical research paradigm, we expect the findings from these studies to have a higher contribution to subsequent clinical research than findings from *in silico*-based and *in vitro*-based studies. A similar or lower contribution from NHP studies would suggest that clinical research is becoming less reliant on this more costly and ethically questionable type of research, thus suggesting that the time for a paradigm shift has come.

## Methods

The design of this study was based on a previously developed method consisting of a quantitative citation analysis and a systematic qualitative analysis of citations.<sup>30</sup>

### Quantitative citation analysis

**Bibliographic search:** The citation analysis was performed between September 2016 and June 2017. The PubMed bibliographic database was searched for papers that described studies employing either NHPs, or *in vitro* or *in silico* research approaches, to investigate MDD. The following Medical Subject Heading (MeSH) search terms were used: ‘Depressive Disorder, Major’ AND MeSH terms: ‘primate’ OR ‘ape’ OR ‘macaque’ OR ‘macaca’ OR ‘rhesus’ OR

'chimpanzee' OR 'bonobo' OR 'gorilla' OR 'gorila' OR 'Pan' OR 'orangutang' OR 'orang-utan' OR 'Orang utan' OR 'orangutan' OR 'ourang-outang' OR 'Pongo' OR 'gibbon' OR 'Hylobates' OR 'Colobus' OR 'Baboon' OR 'Papio' OR 'Mandrillus' OR 'Mandrill' OR 'Cebus' OR 'Cebuella' OR 'Brachyteles' OR 'Loris' OR 'Nycticebus' OR 'lemur' OR 'Callithrix' OR '*in silico*' OR 'computer model' OR 'mathematical model' OR 'computer simulation' OR '*in vitro*' OR 'cell culture' OR 'culture technique' OR 'cell line' OR 'organ culture' OR 'tissue culture'.

MeSH terms are a comprehensive list of key terms related to each human disorder, designed to identify all relevant studies in a given area.<sup>31</sup> Thus, searching for 'Depressive Disorder, Major' retrieves other nomenclatures for the same disorder (e.g. Melancholia). There were no exclusive MeSH terms for NHPs, so the search retrieved additional papers with non-human animals that were excluded by manual sorting. All *in vitro*-based and *in silico*-based papers that used animal data (e.g. rat cell line data) were also excluded.

Papers from scientific journals, books, research reports and conference proceedings written in English, Portuguese or Italian were included (being within the authors' linguistic fluencies). PubMed filters were used, in order to exclude review papers ('review', 'systematic review', 'meta-analysis', 'bibliography'), as well as editorials and other types of non-research papers ('biography', 'auto-biography', 'comment', 'opinion paper', 'interview'), since the aim of the study was to evaluate the impact of original data. The search was restricted to publications prior to 31 December 2011, to allow adequate time for subsequent citation of papers.<sup>32</sup> Nineteen NHP study-based papers, 29 *in silico*-based papers and 38 *in vitro*-based papers describing data from original MDD research were retrieved (see Appendix 1).

**Citation data:** A citation analysis on the retrieved papers was performed by using the cited reference search facility within the Web of Science bibliographic database. For each retrieved paper, the subsequent papers that cited it were identified, and three types of citation data were recorded:

- the total number of times that the retrieved paper was cited;
- the total number of times that the retrieved paper was cited per research category; and
- the total number of times that the retrieved paper was cited per research subject, that is, on MDD or other subjects, as detailed below.

Each citing paper was ascribed to one or more of the following eight research categories: 'invasive animal research'; 'human research'; 'review'; 'opinions' (including editorials, comments or replies to comments); '*in vitro*'; '*in silico*'; 'non-invasive animal research' (e.g.

observational studies with wild animals); and 'other human studies' (e.g. on social perceptions). The term 'human research' referred to any human-based research that might involve, among other things, the analysis of biological samples, epidemiological and behavioural studies, medical case studies and clinical studies. A citing paper could be allocated to more than one category, if it described different research approaches. Whenever the category of the citing paper could not be defined (due to language barriers or absence of an abstract), the paper was labelled as 'not available' and removed from further analysis.

Among the categories 'human research', '*in silico*', '*in vitro*' and 'invasive animal research', it was also recorded whether the citing paper focused on MDD or on other subjects.

**Statistical analysis:** To test for differences between the numbers of citations across research approaches, three generalised linear models (GLMs), each with a Poisson response and a log link function, were implemented. Each model tested one of the following response variables: (a) the total number of citations; (b) the total number of citations by papers in the category 'human research'; and (c) the total number of citations by papers in the category of 'human research' that focused specifically on MDD. In each model, the only explanatory variable was the type of research approach, of which there were three: NHP studies, *in silico*-based approaches and *in vitro*-based approaches. The GLM's goodness of fit was evaluated by visual inspection of the diagnostic plots. Additionally, a Gaussian GLM was used to evaluate whether the proportions of citations by human research papers, and by human research papers specifically on MDD, were different across the three approaches. The analyses were performed in R 3.6.1,<sup>33</sup> by using the function *glm*. The results were considered significant when  $p < 0.05$ .

### Systematic qualitative analysis of citations

Citing papers featuring human research specifically on MDD were systematically analysed by two independent raters, to qualitatively evaluate the contribution of knowledge from NHP studies, or from *in vitro*-based or *in silico*-based research approaches, to the respective human clinical study. Each study was rated according to the following classes, which were defined prospectively, as in Carvalho et al.<sup>30</sup>:

- **Redundant:** when the cited study was only mentioned among other studies as an example. In the case where multiple studies were used as examples of one or more points, the raters were instructed to rate the study as redundant only if there were older or human studies stating exactly the same points.

- *Minor relevance*: when the cited study was cited in either the Discussion or the Introduction, to provide information not directly related to the hypothesis explored in the human study.
- *Relevant to the hypothesis*: when the cited study was cited in the Introduction, to provide information relevant to the hypothesis explored in the human study.
- *Relevant to the methods*: when the human study used the same methodology as that described in the cited paper, with the exception of species differences in the case of NHP study methods.

A paper considered to be ‘relevant’ could be both relevant for the hypothesis and the methods. The other options in the classes are mutually exclusive. In all cases, disagreement between the raters was resolved via detailed discussions until a consensus was reached.

Whenever it was not possible to assess the contribution of a cited paper to a human study due to unavailability of the full publication on the human study, the human research paper was labelled as ‘not available’ and removed from further analysis.

A statistical test was used for comparing proportions (Pearson’s  $\chi^2$  test implemented via R’s *prop.test* function), in order to assess differences between the three cited approaches (i.e. NHP studies, and *in vitro*-based and *in silico*-based approaches). Since, even for the pair with the largest difference, the null hypothesis of equal proportions could not be rejected under the usual significance levels, corrections for multiple comparisons were not attempted.

## Results

### Citation analysis

**NHP study-based results:** Nineteen publications featuring NHP studies in the field of MDD research were retrieved, which were subsequently cited 841 times in total. Of these 19 papers, five featured both human and NHP data.

The subsequent citing papers belonged to the following categories: invasive animal research (312); reviews (245); human research (152); *in vitro* research (81); *in silico* research (14); non-invasive animal research (6); and opinions, including editorials, comments or replies to comments (4). Eighty-five citing papers were not categorised due to being unavailable or written in a language other than English, Portuguese or Italian.

Of the 312 citations by animal research papers, 63 were specifically focused on MDD; of the 152 citations by human research papers, 71 were specifically focused on MDD.

**In silico-based approach results:** Twenty-nine publications describing the use of *in silico*-based approaches in the context of MDD research were retrieved, which were

subsequently cited 806 times in total. Of these 29 papers, seven featured both patient data and computer simulations.

The subsequent citing papers belonged to the following categories: human research (317); *in silico* research (193); reviews (193); invasive animal research (44); *in vitro* research (17); and opinions (17). Fifty-eight citing papers were not categorised due to being unavailable or written in a language other than English, Portuguese or Italian.

Of the 317 citations by human research papers, 94 specifically focused on MDD; of the 193 citations by *in silico*-based research papers, 36 specifically focused on MDD.

**In vitro-based approach results:** Thirty-eight publications describing the use of *in vitro*-based approaches in the context of MDD research were retrieved, which were subsequently cited 2,574 times in total. All of the *in vitro*-based papers used samples of human biological material, mostly being obtained from MDD patients (in 34 out of the 38 studies).

The subsequent citing papers belonged to the following categories: *in vitro* research (1,239), resorting to the use of human biological material (789), laboratory animal biological material (373) or biological material from both sources (12); human research (978), of which 189 studies solely used human participants without concurrent use of *in vitro*-based research approaches; reviews (844); invasive animal research (464), of which 79 studies solely used live animals without concurrent use of *in vitro*-based research approaches; opinions (27); and *in silico* research (16). One hundred and fifty-four citing papers were not categorised due to being unavailable or written in a language other than English, Portuguese or Italian.


Of the 978 citations by human research papers, 482 specifically focused on MDD; of the 1,239 citations by *in vitro* research papers, 487 specifically focused on MDD.

### Comparison of citations of papers based on NHP studies, in vitro approaches and in silico approaches

An inspection of the diagnostic plots showed no reason for concern with regard to the GLM fit. Among the papers using an *in vitro*-based approach, one was frequently cited (711 citations). We performed the analysis both with and without this potential outlier and found no significant differences between the two scenarios.

The GLM estimated the average number of citations per paper for each of the three approaches (Figure 1 (a)). Each NHP paper was cited 3.73 times (standard error (SE): 0.03). Papers based on *in silico* approaches were cited less frequently than this (3.29 times; i.e.  $-0.44$ , SE: 0.05), and papers based on *in vitro* approaches were cited more frequently (4.23 times; i.e.  $+0.5$ , SE: 0.04). Both differences were statistically significant ( $p < 0.0001$ ).

With regard to the average number of subsequent citations by human research papers (Figure 1 (b)), each NHP



**Figure 1.** The number of citations received by the retrieved papers, according to research approach. A bibliographic search was carried out to retrieve papers on MDD, which were categorised as based on NHP studies or *in silico* or *in vitro* approaches, according to the research method described. A citation analysis was then performed to identify papers that subsequently cited these retrieved papers. The graphs show: (a) the total number of times that the retrieved papers were cited, according to their research approach; (b) the number of times that the retrieved papers were cited by papers on human research, according to their research approach; and (c) the number of times that the retrieved papers were cited by papers on human research specifically focused on MDD, according to their research approach. For visualisation purposes, the largest observation in the 'In vitro' category was excluded from the data used to generate the graphs. MDD: major depressive disorder; NHP: non-human primate.

paper was cited 2.03 times (SE: 0.08). In comparison, papers based on *in vitro* and *in silico* approaches were more frequently cited (+1.09, SE: 0.09 and +0.33, SE: 0.10, respectively). These differences were statistically significant ( $p < 0.001$ ).

When looking at the average numbers of citations by human research papers specifically focused on MDD (Figure 1 (c)), each NHP paper was cited 1.27 times (SE: 0.12), which was not statistically different from the number of citations of papers based on *in silico* approaches (−0.12, SE: 0.16). In these MDD-specific publications, papers based on *in vitro* approaches received, on average, more citations (+1.3, SE: 0.13) than papers based on NHP studies, and the difference was statistically significant ( $p < 0.001$ ).

The estimated proportion of citations of NHP papers by human research papers was 0.13 (SE: 0.05). This proportion was significantly higher for papers based on *in silico*

approaches (+0.20, SE: 0.07,  $p = 0.004$ ) and also for papers based on *in vitro* approaches (+0.30, SE: 0.07,  $p < 0.0001$ ).

The estimated proportion of citations of NHP papers by human research papers specifically focused on MDD was 0.06 (SE: 0.03), which was not significantly different from the proportion of citations of papers based on *in silico* approaches (+0.06, SE: 0.04,  $p = 0.1389$ ). The proportion of citations in these MDD-specific publications, of papers based on *in vitro* approaches (+0.14, SE: 0.04), was significantly different from that of the NHP papers ( $p = 0.001$ ).

### Systematic qualitative analysis of citations

Of the 71 human research papers specifically focused on MDD that cited NHP papers, 50 (70%) were fully available for further analysis, along with 401 of the 482 (83%)

**Table 1.** The relevance of cited NHP study-based, *in silico*-based or *in vitro*-based papers to subsequent (i.e. citing) human research papers focused on MDD.<sup>a</sup>

Citations which are:	Papers based on NHP studies	Papers based on <i>in silico</i> approaches	Papers based on <i>in vitro</i> approaches	Total
Redundant or of minor relevance	42 (84%)	43 (75%)	301 (75%)	<b>386</b>
Relevant to the hypothesis or to the methods	8 (16%)	15 (25%)	100 (25%)	<b>123</b>
<b>Total</b>	<b>50</b>	<b>58</b>	<b>401</b>	<b>509</b>

MDD: major depressive disorder; NHP: non-human primate.

<sup>a</sup>The relevance or redundancy of the cited paper to the hypothesis or methods of the citing MDD paper was evaluated by two independent raters. Bold: total value.

human research papers on MDD that cited *in vitro*-based papers, and 58 of the 94 (62%) human research papers on MDD that cited *in silico*-based papers. It was judged that eight of 50 (16%), 15 of 58 (25%) and 100 of 401 (25%) of citations of papers based on NHP studies, *in silico* and *in vitro* approaches, respectively, were relevant to the hypothesis and/or the methods in the citing human research paper on MDD (see Table 1).

The statistical test used to compare the proportions did not reveal any significant differences between the proportions of relevant citations between NHP–*in vitro*, NHP–*in silico* and *in vitro*–*in silico* ( $p = 0.31, 0.20$  and  $1$ , respectively).

## Discussion

We quantitatively and qualitatively analysed the contribution of NHP, *in vitro* and *in silico*-based research approaches to the contemporary understanding of MDD. Of the three approaches analysed, NHP studies seemed to be the approach that was least likely to contribute to furthering progress in this field of human medical research. Of the three, the human-based *in vitro* approach seemed to influence human research to the greatest extent, judging by the number of citations. However, all three approaches seemed to be equally relevant in informing the hypothesis and/or methods of subsequent human research studies.

Overall, our results suggest that these less funded non-animal research approaches<sup>34</sup> are more or equally effective than heavily invested animal-based research in reaching their final goal — which is to inform clinical research to improve human healthcare. Our quantitative results showed that *in silico*-based and *in vitro*-based approaches contributed more than NHP study-based approaches to human medical research, as the proportion of cited papers featuring the former two approaches was higher than the proportion of cited papers featuring the latter. NHP study-based papers were mainly cited by other papers on animal experimentation, which suggests that they are mainly contributing to subsequent animal research rather than to advances in human healthcare. *In vitro* studies seemed to be the most effective approach, since this approach received significantly more citations in total, and by human research

papers either specifically focused on MDD or on other general medical areas.

Of the five analysed NHP study-based papers that were relevant to the citing human research papers on MDD in terms of their hypothesis, method or both, one featured both NHP and human research data. This paper was cited twice, and both citing papers referred to the human research data rather than to the NHP data. Another one of these five NHP papers was considered relevant to the methods and was cited once. The citing paper described both human and rhesus monkey data, and the citation was relevant to the methods used with the rhesus monkeys. After excluding these cases, only three out of the 19 NHP studies were relevant to the hypothesis and/or methods of the subsequent human research studies on MDD.

The results of our citation analysis also suggest that the widely accepted approach to testing medical hypotheses — which relies on *in vitro*-based and *in silico*-based research as a preliminary step prior to animal testing — is not actually working as intended, since clinical papers tend to cite *in silico*-based and *in vitro*-based papers directly too. However, citations of *in silico*-based and *in vitro*-based papers in subsequent publications on human clinical studies of MDD constituted a low percentage (50% or less) of the total citations received in all three analysed categories. This may be explained by the complexity of MDD, which shares certain genetic factors, phenotypic traits and possible neurologic pathways with a number of other disorders. Hence, a human study on anorexia might cite a non-clinical study on MDD focused on weight loss, since weight change is one of the symptoms of MDD.

As to the qualitative results, the judged relevance of the initially retrieved papers to the publications subsequently citing them was low for all three analysed research approaches. Even though a higher percentage of cited *in silico*-based and *in vitro*-based papers were relevant to the hypothesis and/or methods used by the citing clinical studies, the differences between the three approaches, in the extent of their judged relevance, were deemed insignificant. However, the size of the observed effect — where the proportion of citations of NHP-based papers was much lower than that of *in silico*-based or *in vitro*-based papers — suggests that, while not statistically

significant (due to lack of statistical power), there might be a relevant practical difference.

Several important developments in *in vitro* technologies (e.g. organs-on-chips<sup>35</sup>) and in *in silico* technologies (e.g. advanced artificial intelligence based on sophisticated machine learning tools<sup>36</sup>) have been published since 2011. Such studies have been excluded from our analysis, in order to ensure that sufficient time is given to allow for subsequent citation of the resulting papers. However, it is reasonable to expect that these cutting-edge technologies are currently being widely used to generate and test new hypotheses in human medicine.<sup>27</sup> Similarly, induced pluripotent stem cells, even though they have been worked on and developed for more than a decade,<sup>37</sup> have only recently been recommended for MDD research.<sup>38</sup> In light of the above, it would be interesting to repeat the current study a decade from now to investigate whether this has led to an increase in the number of subsequent citations of *in vitro*-based and *in silico*-based papers on MDD, in both MDD-focused and general human research publications.

We recognise that our study has certain limitations. Due to resource constraints, we were unable to use a greater number of search engines (e.g. CAB Abstracts). This would have increased the likelihood of retrieving all *in silico*-based, *in vitro*-based and NHP study-based papers on MDD, which would have increased our sample size and thus made it more comprehensive. Similarly, we were unable to examine the reference lists of many of the retrieved papers, in order to locate additional relevant papers. This inevitably means that some relevant publications might not have been identified. Because the sample size was small, our results should be interpreted with this caveat.

Finally, we are aware of the difficulty in objectively determining the relevance of a cited paper to the publication citing it. We used two different raters, in order to attempt to decrease any error in subjective assessment. Occasionally, the raters differed in their initial assessment, indicating that, even when the same criteria are used for assessment, differences can sometimes arise. However, our experience suggests that these differences would relate to only a small proportion of the papers assessed. Despite the limitations in the citation search and in the systematic qualitative analysis of the citation value, we consider that the method we followed is useful when evaluating the effectiveness of different research approaches. We hope that similar studies adopt this methodology, in order to investigate other medical disorders.

Our results suggest that the contribution of NHP studies to the current understanding of MDD is poor, and that other approaches with potentially superior relevance to humans should be used. Our results also shed light on the controversy around the efficacy of NHP-based research for investigating human disorders. This controversy is longstanding, with some authors claiming that their use is

crucial for medical advancement,<sup>23</sup> while others assert the opposite.<sup>39,40</sup> However, ongoing scientific advances in non-animal methods for the acquisition of knowledge and the development of new treatments may provide future alternative solutions to help avoid the dilemmas and concerns surrounding NHP use.

## Conclusions

To our knowledge, this is the first study to compare the effectiveness of original studies involving the use of NHP, *in vitro* and *in silico* research approaches to inform the medical research community within the MDD field. Our results suggest that, in this field of medical research, human-based *in vitro* and *in silico* research approaches are more promising than NHP studies, in generating new hypotheses and methods for subsequent clinical research.

Given the scientific advances in human-based research methods, we suggest that our methodology could be used in the future to analyse the impact of more recent technologies in informing human medical research. Such analysis could examine if and how the standard paradigm for testing medical hypotheses is still being followed, from applied research, through animal use in preclinical testing, and on to clinical research and development. It could also provide further insight into how the 'gold standard' that considers *in vitro*-based and *in silico*-based research approaches as merely preliminary steps prior to animal testing could be challenged and revised. Given the scientific and ethical solutions that innovative human-based approaches are providing, with relatively little investment when compared to the investments in animal-based research, a reallocation of resources is clearly warranted in favour of researching and developing the use of such approaches as part of human medical research.

## Acknowledgements

The authors would like to acknowledge the University of Lisbon EcoComp team members for their valuable insights during the conception and design of this study.

## Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship and/or publication of this article.

## Funding

The author(s) disclosed receipt of the following financial support for the research, authorship and/or publication of this article: This study was financed by Animalfree Research–Switzerland, and by Portuguese national funds through FCT–Fundação para a Ciência e a Tecnologia, within the CFCUL Unit funding UID/FIL/00678/2013. TAM received partial support from CEAUL (funded by FCT–Fundação para a Ciência e a Tecnologia, Portugal, through the project UID/MAT/00006/2019). Open access publication costs were covered by Animalfree Research–Switzerland and by

funding provided by the Ketty and Leif Hjordt Foundation. None of these funders played any role in the study conceptualisation, design, conduct, data collection or analysis, authorship of the resulting manuscript or the decision to submit it for publication.

## Ethical approval

Ethics approval was not required for this article.

## Informed consent

Informed consent was not required for this article.

## References

- WHO. *Depression, key facts*. Geneva: World Health Organization, <https://www.who.int/news-room/fact-sheets/detail/depression> (2018, accessed 29 June 2019).
- Cuijpers P, Vogelzangs N, Twisk J, et al. Comprehensive meta-analysis of excess mortality in depression in the general community versus patients with specific illnesses. *Am J Psychiatry* 2014; 171: 453–462.
- Li CT, Bai YM, Tu PC, et al. Major depressive disorder and stroke risks: a 9-year follow-up population-based, matched cohort study. *PLoS One* 2012; 7: e46818.
- Rauma PH, Pasco JA, Berk M, et al. The association between major depressive disorder, use of antidepressants and bone mineral density (BMD) in men. *J Musculoskelet Neuronal Interact* 2015; 15: 177–185.
- American Psychiatric Association. *DSM-5: Manual diagnóstico e estatístico de transtornos mentais*. São Paulo: Artmed Editora, 2014, p. 992.
- Vos T, Allen C, Arora M, et al. Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 2016; 388: 1545–1602.
- Sobocki P, Jonsson B, Angst J, et al. Cost of depression in Europe. *J Ment Health Policy Econ* 2006; 9: 87–98.
- Wittchen HU, Jacobi F, Rehm J, et al. The size and burden of mental disorders and other disorders of the brain in Europe 2010. *Eur Neuropsychopharmacol* 2011; 21: 655–679.
- Greenberg PE, Fournier AA, Sisitsky T, et al. The economic burden of adults with major depressive disorder in the United States (2005 and 2010). *J Clin Psychiatry* 2015; 76: 155–162.
- Maurer K and Quimby F. Animal models in biomedical research. In: Fox JG, Anderson LC, Otto GM, Pritchett-Corning KR and Whary MT (eds) *Laboratory animal medicine*. 4th ed. Cambridge: Academic Press, 2015, pp. 1497–1534.
- Anon. *Special Eurobarometer 340/Wave 73.1: science and technology*. Report, European Commission, Brussels, June 2010, [http://ec.europa.eu/commfrontoffice/publicopinion/archives/ebs/ebs\\_340\\_en.pdf](http://ec.europa.eu/commfrontoffice/publicopinion/archives/ebs/ebs_340_en.pdf) (accessed 29 June 2019).
- Lankau EW, Turner PV, Mullan RJ, et al. Use of nonhuman primates in research in North America. *J Am Assoc Lab Anim Sci* 2014; 53: 278–282.
- Anon. *Animal Welfare Act 1999*. Wellington: New Zealand Government, <http://www.legislation.govt.nz/act/public/1999/0142/59.0/DLM49664.html> (1999, accessed 19 August 2019).
- Kraskov A, Prabhu G, Quallo MM, et al. Ventral premotor–motor cortex interactions in the macaque monkey during grasp: response of single neurons to intracortical microstimulation. *J Neurosci* 2011; 31: 8812–8821.
- Vigneswaran G, Philipp R, Lemon RN, et al. M1 corticospinal mirror neurons and their role in movement suppression during action observation. *Curr Biol* 2013; 23: 236–243.
- Maninger N, Capitanio JP, Mason WA, et al. Acute and chronic stress increase DHEAS concentrations in rhesus monkeys. *Psychoneuroendocrinology* 2010; 35: 1055–1062.
- Knight A. Laboratory animal statistics for Great Britain: implications for animal welfare. *AATEX* 2010; 17: 51–52.
- Gottlieb DH, Capitanio JP and McCowan B. Risk factors for stereotypic behavior and self-biting in rhesus macaques (*Macaca mulatta*): animal's history, current environment, and personality. *Am J Primatol* 2013; 75: 995–1008.
- Morgan KN and Tromborg CT. Sources of stress in captivity. *Appl Anim Behav Sci* 2007; 102: 262–302.
- Buller T. Animal minds and neuroimaging: bridging the gap between science and ethics? *Camb Q of Health Ethics* 2014; 23: 173–181.
- Keen J. Wasted money in United States biomedical and agricultural animal research. In: Herrmann K and Jayne K (eds) *Animal experimentation: working towards a paradigm change*. Leiden: Brill, 2019, pp. 244–272.
- Fitzgerald TA. Comparison of research cost: man–primate animal–other animal models. *J Med Primatol* 1982; 12: 138–145.
- Worlein JM. Nonhuman primate models of depression: effects of early experience and stress. *ILAR J* 2014; 55: 259–273.
- Pound P and Bracken MB. Is animal research sufficiently evidence based to be a cornerstone of biomedical research? *BMJ* 2014; 348: g3387.
- Carvalho C, Alves D, Knight A, et al. Is animal-based biomedical research being used in its original context? In: Herrmann K and Jayne K (eds) *Animal experimentation: working towards a paradigm change*. Leiden: Brill, 2019, pp. 376–390.
- Benam KH, Gilchrist S, Kleensang A, et al. Exploring new technologies in biomedical research. *Drug Discov Today* 2019; 24: 1242–1247.
- Archibald K, Tsaioun K, Kenna JG, et al. Better science for safer medicines: the human imperative. *J R Soc Med* 2018; 111: 433–438.
- Taylor K. Recent developments in alternatives to animal testing. In: Herrmann K and Jayne K (eds) *Animal experimentation: working towards a paradigm change*. Leiden: Brill, 2019, pp. 583–609.
- Langley GR, Adcock IM, Busquet F, et al. Towards a 21st-century roadmap for biomedical research and drug discovery: consensus report and recommendations. *Drug Discov Today* 2017; 22: 327–339.



30. Carvalho C, Crespo MV, Bastos LF, et al. Contribution of animal models to contemporary understanding of attention deficit hyperactivity disorder. *ALTEX* 2016; 33: 243–249.
31. Uman L. Systematic reviews and meta-analyses. *J Can Acad Child Adolesc Psychiatry* 2011; 20: 57–59.
32. Wooding S, Pollitt A, Castle-Clarke S, et al. Mental Health Retrosight: Understanding the returns from research (lessons from schizophrenia): policy report. *Rand Health Q* 2014; 4(1): 8.
33. R Core Team. *R: a language and environment for statistical computing*. Vienna: R Foundation for Statistical Computing, <https://www.R-project.org> (2018, accessed 21 August 2019).
34. Frank J. Technological lock-in, positive institutional feedback, and research on laboratory animals. *Struct Change Econ D* 2005; 16: 557–575.
35. Bhatia SN and Ingber DE. Microfluidic organs-on-chips. *Nature Biotech* 2014; 32: 760–772.
36. Batool M, Ahmad B and Choi S. A structure-based drug discovery paradigm. *Int J Mol Sci* 2019; 20: 2783.
37. Romito A and Cobellis G. Pluripotent stem cells: current understanding and future directions. *Stem Cells Int* 2016; 2016: 9451492.
38. Licinio J and Wong ML. Serotonergic neurons derived from induced pluripotent stem cells (iPSCs): a new pathway for research on the biology and pharmacology of major depression. *Mol Psychiatry* 2016; 21: 1–2.
39. Knight A. The poor contribution of chimpanzee experiments to biomedical progress. *J Appl Anim Welf Sci* 2007; 10: 281–308.
40. Bailey J and Taylor K. Non-human primates in neuroscience research: the case against its scientific necessity. *Altern Lab Anim* 2016; 44: 43–69.

## Appendix I: A list of retrieved articles

### In vitro-based approach

Bartlett JA, Demetrikopoulos MK, Schleifer SJ, et al. Phagocytosis and killing of *Staphylococcus aureus*: effects of stress and depression in children. *Clin Diagn Lab Immunol* 1997; 4: 362–366.

Bertilsson L, Aberg-Wistedt A, Lidén A, et al. Alprazolam does not inhibit the metabolism of nortriptyline in depressed patients or inhibit the metabolism of desipramine in human liver microsomes. *Ther Drug Monit* 1988; 10: 231–233.

Brusov OS, Beliaev BS, Katasonov AB, et al. Development of subsensitivity to imipramine in the system of reverse serotonin uptake by thrombocytes in patients with endogenous depression. *Zh Nevropatol Psikiatr Im S S Korsakova* 1988; 88: 96–100.

Chen B, Dowlatshahi D, MacQueen GM, et al. Increased hippocampal BDNF immunoreactivity in subjects treated with antidepressant medication. *Biol Psychiatry* 2001; 50: 260–265.

Conklin SM, Runyan CA, Leonard S, et al. Age-related changes of n-3 and n-6 polyunsaturated fatty acids in the anterior cingulate cortex of individuals with major depressive disorder. *Prostaglandins Leukot Essent Fatty Acids* 2010; 82: 111–119.

De Paermentier F, Cheetham SC, Crompton MR, et al. Lower cortical beta-adrenoceptor binding sites in post-mortem samples from depressed suicide victims. *Br J Pharmacol* 1989; 98: 818P.

Du L, Faludi G, Palkovits M, et al. Frequency of long allele in serotonin transporter gene is increased in depressed suicide victims. *Biol Psychiatry* 1999; 46: 196–201.

Ebstein RP, Lerer B, Shapira B, et al. Cyclic AMP second-messenger signal amplification in depression. *Br J Psychiatry* 1988; 152: 665–669.

Franke L, Schewe HJ, Uebelhack R, et al. Platelet-5HT uptake and gastrointestinal symptoms in patients suffering from major depression. *Life Sci* 2003; 74: 521–531.

Garbin L, Bianchin GL, De Bertolini C, et al. Inhibitory effect of imipramine on epinephrine-dependent platelet aggregation: “in vitro” studies on platelets from healthy and depressed people. *Pharmacol Res Commun* 1983; 15: 23–27.

Gilad GM, Gilad VH, Casanova MF, et al. Polyamines and their metabolizing enzymes in human frontal cortex and hippocampus: preliminary measurements in affective disorders. *Biol Psychiatry* 1995; 38: 227–234.

Greenstein AS, Paranthaman R, Burns A, et al. Cerebrovascular damage in late-life depression is associated with structural and functional abnormalities of subcutaneous small arteries. *Hypertension* 2010; 56: 734–740.

Heiser P, Lanquillon S, Krieg JC, et al. Differential modulation of cytokine production in major depressive disorder by cortisol and dexamethasone. *Eur Neuropsychopharmacol* 2008; 18: 860–870.

Himmerich H, Fulda S, Sheldrick AJ, et al. IFN- $\gamma$  reduction by tricyclic antidepressants. *Int J Psychiatry Med* 2010; 40: 413–424.

Hing B, Davidson S, Lear M, et al. A polymorphism associated with depressive disorders differentially regulates brain derived neurotrophic factor promoter IV activity. *Biol Psychiatry* 2012; 71: 618–626.

- Hrdina PD, Demeter E, Vu TB, et al. 5-HT uptake sites and 5-HT<sub>2</sub> receptors in brain of antidepressant-free suicide victims/depressives: increase in 5-HT<sub>2</sub> sites in cortex and amygdala. *Brain Res* 1993; 614: 37–44.
- Irwin M, Costlow C, Williams H, et al. Cellular immunity to varicella-zoster virus in patients with major depression. *J Infect Dis* 1998; 178: S104–S108.
- Karege F, Bovier P, Widmer J, et al. Decrease in epinephrine-induced attenuation of platelet adenylate cyclase activity in depressed patients: relation with plasma electrolytes. *Neuropsychobiology* 1992; 26: 129–135.
- Kim YK, Lee SW, Kim SH, et al. Differences in cytokines between non-suicidal patients and suicidal patients in major depression. *Prog Neuropsychopharmacol Biol Psychiatry* 2008; 32: 356–361.
- Klimek V, Roberson G, Stockmeier CA et al. Serotonin transporter and MAO-B levels in monoamine nuclei of the human brainstem are normal in major depression. *J Psychiatr Res* 2003; 37: 387–397.
- Kok FW, Heijnen CJ, Bruijn JA, et al. Immunoglobulin production *in vitro* in major depression: a pilot study on the modulating action of endogenous cortisol. *Biol Psychiatry* 1995; 38: 217–226.
- Kubera M, Kenis G, Bosmans E, et al. Stimulatory effect of antidepressants on the production of IL-6. *Int Immunopharmacol* 2004; 4: 185–192.
- Lawrence KM, De Paermentier F, Cheetham SC, et al. Brain 5-HT uptake sites, labelled with [<sup>3</sup>H]-paroxetine, in post-mortem samples from depressed suicide victims. *Br J Pharmacol* 1989; 98: 812P.
- Lawrence KM, Falkowski J, Jacobson RR, et al. Platelet 5-HT uptake sites in depression: three concurrent measures using [<sup>3</sup>H] imipramine and [<sup>3</sup>H] paroxetine. *Psychopharmacol* 1993; 110: 235–239.
- Lin A, Song C, Kenis G, et al. The *in vitro* immunosuppressive effects of moclobemide in healthy volunteers. *J Affect Disord* 2000; 58: 69–74.
- Maes M and Thompson P. Analysis of partial variance to control for day-to-day variability in functional immune tests in depression. *Neuropsychobiology* 1997; 36: 107–111.
- Mann JJ, Brown RP, Halper JP, et al. Reduced sensitivity of lymphocyte beta-adrenergic receptors in patients with endogenous depression and psychomotor agitation. *N Eng J Med* 1985; 313: 715–720.
- Miguel-Hidalgo JJ, Baucom C, Dilley G, et al. Glial fibrillary acidic protein immunoreactivity in the prefrontal cortex distinguishes younger from older adults in major depressive disorder. *Biol Psychiatry* 2000; 48: 861–873.
- Mikuni M, Kusumi I, Kagaya A, et al. Increased 5-HT<sub>2</sub> receptor function as measured by serotonin-stimulated phosphoinositide hydrolysis in platelets of depressed patients. *Prog Neuropsychopharmacol Biol Psychiatry* 1991; 15: 49–61.
- Miller GE, Rohleder N, Stetler C, et al. Clinical depression and regulation of the inflammatory response during acute stress. *Psychosom Med* 2005; 67: 679–687.
- Mizrahi C, Stojanovic A, Urbina M, et al. Differential cAMP levels and serotonin effects in blood peripheral mononuclear cells and lymphocytes from major depression patients. *Int Immunopharmacol* 2004; 4: 1125–1133.
- Molnar M, Potkin SG, Bunney WE, et al. MRNA expression patterns and distribution of white matter neurons in dorsolateral prefrontal cortex of depressed patients differ from those in schizophrenia patients. *Biol Psychiatry* 2003; 53: 39–47.
- Morishita S, Aoki S and Watanabe S. Different effect of desipramine on protein kinase C in platelets between bipolar and major depressive disorders. *Psychiatry Clin Neurosci* 1999; 53: 11–15.
- Perry EK, Marshall EF, Blessed G, et al. Decreased imipramine binding in the brains of patients with depressive illness. *Br J Psychiatry* 1983; 142: 188–192.
- Szentistvanyi I, Janka Z and Rimanoczy A. Alteration of erythrocyte phosphate transport in primary depressive disorders. *J Affect Disord* 1980; 2: 229–238.
- Valdizán EM, Gutierrez O and Pazos A. Adenylate cyclase activity in postmortem brain of suicide subjects: reduced response to  $\beta$ -adrenergic stimulation. *Biol Psychiatry* 2003; 54: 1457–1464.
- Wood PL, Suranyi-Cadotte BE, Nair NPV, et al. Lack of association between [<sup>3</sup>H]imipramine binding sites and uptake of serotonin in control, depressed and schizophrenic patients. *Neuropharmacol* 1983; 22: 1211–1214.

### In silico-based approach

- Becker S, MacQueen G and Wojtowicz JM. Computational modeling and empirical studies of hippocampal neurogenesis-dependent memory: effects of interference, stress and depression. *Brain Res* 2009; 1299: 45–54.
- Bulmash EL, Moller HJ, Kayumov L, et al. Psychomotor disturbance in depression: assessment using a driving simulator paradigm. *J Affect Disord* 2006; 93: 213–218.
- Bursi R, Erdemli G, Campbell R, et al. Translational PK–PD modelling of molecular target modulation for the AMPA receptor positive allosteric modulator Org 26576. *Psychopharmacol* 2011; 218: 713–724.

Costafreda SG. Parametric coordinate-based meta-analysis: valid effect size meta-analysis of studies with differing statistical thresholds. *J Neurosci Methods* 2012; 210: 291–300.

Demirtas H and Doganay B. Simultaneous generation of binary and normal data with specified marginal and association structures. *J Biopharm Stat* 2012; 22: 223–236.

Fischer K, Goetghebuer E, Vrijens B, et al. A structural mean model to allow for noncompliance in a randomized trial comparing 2 active treatments. *Biostatistics* 2010; 12: 247–257.

Gardner W, Shear K, Kelleher KJ, et al. Computerized adaptive measurement of depression: a simulation study. *BMC Psychiatry* 2004; 4: 13.

Giuffra LA and Risch N. Diminished recall and the cohort effect of major depression: a simulation study. *Psychol Med* 1994; 24: 375–383.

Heo M and Leon AC. Sample sizes required to detect two-way and three-way interactions involving slope differences in mixed-effects linear models. *J Biopharm Stat* 2010; 20: 787–802.

Hida E and Tango T. On the three-arm non-inferiority trial including a placebo with a prespecified margin. *Stat Med* 2010; 30: 224–231.

Javaras KN, Hudson JI and Laird NM. Fitting ACE structural equation models to case-control family data. *Genet Epidemiol* 2009; 34: 238–245.

Koshelev M, Lohrenz T, Vannucci M, et al. Biosensor approach to psychopathology classification. *PLoS Comput Biol* 2010; 6: e1000966.

Merlo-Pich E, Alexander RC, Fava M, et al. A new population-enrichment strategy to improve efficiency of placebo-controlled clinical trials of antidepressant drugs. *Clin Pharmacol Ther* 2010; 88: 634–642.

Partonen T, Treutlein J, Alpmann A, et al. Three circadian clock genes *Per2*, *Arntl*, and *Npas2* contribute to winter depression. *Ann Med* 2007; 39: 229–238.

Patten SB. An animated depiction of major depression epidemiology. *BMC Psychiatry* 2007; 7: 23.

Patten SB. The National Population Health Survey's assessment of depression risk factor associations: a simulation study assessing vulnerability to bias. *Chronic Dis Inj Can* 2012; 32: 70–75.

Santen G, Horrigan J, Danhof M, et al. From trial and error to trial simulation. Part 2: an appraisal of current beliefs in the design and analysis of clinical trials for antidepressant drugs. *Clin Pharmacol Ther* 2009; 86: 255–262.

Serretti A, Olgiaiti P, Bajo E, et al. A model to incorporate genetic testing (5-HTTLPR) in pharmacological treatment of major depressive disorders. *World J Biol Psychiatry* 2011; 12: 501–515.

Shang EY, Gibbs MA, Landen JW, et al. Evaluation of structural models to describe the effect of placebo upon the time course of major depressive disorder. *J Pharmacokinet Pharmacodyn* 2009; 36: 63–80.

Shen J, Moller HJ, Wang X, et al. Mirtazapine, a sedating antidepressant, and improved driving safety in patients with major depressive disorder: a prospective, randomized trial of 28 patients. *J Clin Psychiatry* 2009; 70: 370–377.

Shults J, Sun W, Tu X, et al. A comparison of several approaches for choosing between working correlation structures in generalized estimating equation analysis of longitudinal binary data. *Stat Med* 2009; 28: 2338–2355.

Slade T. Taxometric investigation of depression: evidence of consistent latent structure across clinical and community samples. *Aust N Z J Psychiatry* 2007; 41: 403–410.

Soeteman DI, Miller M and Kim JJ. Modeling the risks and benefits of depression treatment for children and young adults. *Value Health* 2012; 15: 724–729.

Suckling J and Bullmore E. Permutation tests for factorially designed neuroimaging experiments. *Human Brain Mapp* 2004; 22: 193–205.

Taub NA, Morgan Z, Brugha TS, et al. Recalibration methods to enhance information on prevalence rates from large mental health surveys. *Int J Methods Psychiatr Res* 2005; 14: 3–13.

Tretter F, Gebicke-Haerter PJ, An Der Heiden U, et al. Affective disorders as complex dynamic diseases — a perspective from systems biology. *Pharmacopsychiatry* 2011; 44: S2–S8.

Wang X, Lin Y, Song C, et al. Detecting disease-associated genes with confounding variable adjustment and the impact on genomic meta-analysis: with application to major depressive disorder. *BMC Bioinformatics* 2012; 13: 52.

Weigelt K, Carvalho LA, Drexhage RC, et al. TREM-1 and DAP12 expression in monocytes of patients with severe psychiatric disorders. EGR3, ATF3 and PU.1 as important transcription factors. *Brain Behav Immun* 2011; 25: 1162–1169.

Zhang WR, Pandurangi AK and Peace KE. YinYang dynamic neurobiological modeling and diagnostic analysis of major depressive and bipolar disorders. *IEEE Trans Biomed Eng* 2007; 54: 1729–1739.

Zubenko GS and Hughes HB. Effects of the G(-656)A variant on *CREB1* promoter activity in a glial cell line: interactions with gonadal steroids and stress. *Am J Med Genet B Neuropsychiatr Genet* 2008; 147B: 579–585.

### Non-human primate study-based

Eggan SM, Stoyak SR, Verrico CD, et al. Cannabinoid CB1 receptor immunoreactivity in the prefrontal cortex: comparison of schizophrenia and major depressive disorder. *Neuropsychopharmacol* 2010; 35: 2060–2071.

Feyissa AM, Woolverton WL, Miguel-Hidalgo JJ, et al. Elevated level of metabotropic glutamate receptor 2/3 in the prefrontal cortex in major depression. *Prog Neuropsychopharmacol Biol Psychiatry* 2010; 34: 279–283.

Freedman LJ, Insel TR and Smith Y. Subcortical projections of area 25 (subgenual cortex) of the macaque monkey. *J Comp Neurol* 2000; 421: 172–188.

Fuchs E, Czéh B, Michaelis T, et al. Synaptic plasticity and tianeptine: structural regulation. *Eur Psychiatry* 2002; 17(Suppl 3): 311–317.

Goncharova ND, Marenin VY and Oganyan TE. Aging of the hypothalamic-pituitary-adrenal axis in nonhuman primates with depression-like and aggressive behavior. *Aging* 2010; 2: 854–866.

Kraemer GW, Ebert MH, Lake CR, et al. Cerebrospinal fluid measures of neurotransmitter changes associated with pharmacological alteration of the despair response to social separation in rhesus monkeys. *Psychiatry Res* 1984; 11: 303–315.

Kraemer GW and McKinney WT. Interactions of pharmacological agents which alter biogenic amine metabolism and depression: an analysis of contributing factors within a primate model of depression. *J Affect Disord* 1979; 1: 33–54.

Lu NZ, Eshleman AJ, Janowsky A, et al. Ovarian steroid regulation of serotonin reuptake transporter (SERT) binding, distribution, and function in female macaques. *Mol Psychiatry* 2003; 8: 353–360.

Lyons DM, Wang OJ, Lindley SE, et al. Separation induced changes in squirrel monkey hypothalamic-pituitary-adrenal physiology resemble aspects of hypercortisolism in humans. *Psychoneuroendocrinology* 1999; 24: 131–142.

Moscip TD, Terrace HS, Sackeim HA, et al. A primate model of anterograde and retrograde amnesia produced by convulsive treatment. *J ECT* 2004; 20: 26–36.

Nabulsi N, Huang Y, Weinzimmer D, et al. High-resolution imaging of brain 5-HT<sub>1B</sub> receptors in the rhesus monkey using [<sup>11</sup>C]P943. *Nucl Med Biol* 2010; 37: 205–214.

Pickar D, Naber D, Post RM, et al. Measurement of endorphins in CSF. Relationship to psychiatric diagnosis. *Mod Probl Pharmacopsychiatry* 1981; 17: 246–262.

Pongrac JL, Middleton FA, Peng L, et al. Heat shock protein 12A shows reduced expression in the prefrontal cortex of subjects with schizophrenia. *Biol Psychiatry* 2004; 56: 943–950.

Qiao M, Zhao Q, Zhang H, et al. Isolating with physical restraint low status female monkeys during luteal phase might make an appropriate premenstrual depression syndrome model. *J Affect Disord* 2007; 102: 81–91.

Rasmussen KL and Reite M. Loss-induced depression in an adult macaque monkey. *Am J Psychiatry* 1982; 139: 679–681.

Shively CA, Friedman DP, Gage HD, et al. Behavioral depression and positron emission tomography-determined serotonin 1A receptor binding potential in cynomolgus monkeys. *Arch Gen Psychiatry* 2006; 63: 396–403.

Shively CA, Register TC, Adams MR, et al. Depressive behavior and coronary artery atherogenesis in adult female cynomolgus monkeys. *Psychosom Med* 2008; 70: 637–645.

Spellman T, Peterchev AV and Lisanby SH. Focal electrically administered seizure therapy: a novel form of ECT illustrates the roles of current directionality, polarity, and electrode configuration in seizure induction. *Neuropsychopharmacol* 2009; 34: 2002–2010.

Szewczyk B, Albert PR, Rogaeva A, et al. Decreased expression of Freud-1/CC2D1A, a transcriptional repressor of the 5-HT<sub>1A</sub> receptor, in the prefrontal cortex of subjects with major depression. *Int J Neuropsychopharmacol* 2010; 13: 1089–1101.

Van Kampen M, Kramer M, Hiemke C, et al. The chronic psychosocial stress paradigm in male tree shrews: evaluation of a novel animal model for depressive disorders. *Stress* 2002; 5: 37–46.